

SHINGLES (ZOSTER)

Shingles (zoster) is caused by the varicella-zoster virus that remains inactive (dormant) in the body of people who have had chickenpox (varicella). You get shingles from your own chickenpox virus, not from someone else.

This usually occurs when the immune system is weakened for various reasons, including certain illnesses, conditions, treatments, or aging. Although shingles usually occurs in adulthood, children who were infected with varicella in utero or during infancy may develop shingles during childhood.

CAUSE

Varicella-zoster virus, a member of the herpesvirus family.

SYMPTOMS

Severe pain, itching, and numbness along certain nerve pathways, commonly involving one side of the body. About 1 to 3 days later, a red rash appears at the site. Clusters of blisters appear soon after, usually on one side of the body and closer together than in chickenpox. The blisters dry out and crust over within a few days. The rash and pain usually disappear within 3 to 5 weeks. Shingles is a milder illness in children than in adults, but it can be a serious illness in those who have weakened immune systems.

SPREAD

A person must have already had chickenpox disease in the past to develop shingles. Shingles does not spread from one person to another as shingles. When people who have not had chickenpox have contact with the fluid from the shingles blisters, they can develop chickenpox.

INCUBATION

None.

CONTAGIOUS PERIOD

In an otherwise healthy person, from the start of the rash until all the blisters have crusted.

DIAGNOSIS

Providers usually diagnose shingles by observing of the telltale rash. Providers may also take a tissue sample or culture of the blisters to send to the lab.

EXCLUSION

Childcare and School: None if blisters can be completely covered by clothing or a bandage. If blisters cannot be covered, exclude until the blisters have crusted. Persons with severe, disseminated shingles should be excluded regardless of whether the sores can be covered.

TREATMENT

Antiviral medications may be prescribed. DO NOT GIVE ASPIRIN or OTHER SALICYLATE CONTAINING MEDICATIONS TO ANY CHILD OR ADOLESCENT UNDER 18 YEARS OF AGE.

PREVENTION/CONTROL

- Pregnant women or persons with weakened immune systems who have not had chickenpox and are exposed to shingles virus should call their healthcare provider immediately for possible treatment.
- Wash hands thoroughly with soap and warm running water after contact with fluid from blisters or sores.
- Thorough hand washing is the best way to prevent the spread of communicable diseases.
- Susceptible persons (those who have not had chickenpox or varicella vaccine in the past) who have been exposed to someone with shingles should call their healthcare provider immediately. Getting varicella vaccine within 3 days, and possibly up to 5 days, of exposure may prevent disease in these people.
- Clean and disinfect all objects and surfaces contaminated with blister fluid at least daily and when soiled. (See IV-4 through IV-6)

For more information, please contact your local health department.

[Carson City Health & Human Services](#) (Carson City, Douglas, Lyon Counties) 775-434-1690 & 775-887-2190 (after hours)

[Central Nevada Health District](#) (Churchill, Mineral, Eureka, Pershing Counties) 775-866-7535

[Northern Nevada Public Health](#) (Washoe County) 775-328-2447 (24 hours)

[Southern Nevada Health District](#) (Clark County) 775-759-1300 (24 hours)

[Division of Public and Behavioral Health](#) (All other Counties) 775-400-0333 (24 hours)

Email: dpbhepi@health.nv.gov